




## Stilbenes from *Paphiopedilum exul* roots

Naphatsawan P., Nonthalert L., Rutt S. and Suchada S.\*

Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand

**Keywords:** *Paphiopedilum exul*; Orchidaceae; *trans*-stilbenes

Five *trans*-stilbenes including (*E*)-3-methoxy-5-hydroxystilbene (**1**), (*E*)-3,5'-dimethoxy-5,2'-dihydroxystilbene (**2**), thunalbene (**3**), (*E*)-3-methoxy-5,2'-dihydroxystilbene (**4**) and (*E*)-3-methoxy-2-(4-hydroxybenzyl)-5,2'-dihydroxystilbene (**5**) were isolated from the roots of *Paphiopedilum exul* (family Orchidaceae). Their chemical structures were identified by spectroscopic methods including <sup>1</sup>H-NMR, <sup>13</sup>C-NMR and 2D-NMR. This study is the first report of chemical constituents of this lady's slipper orchid native to the south of Thailand.



\* Corresponding author: Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand; Tel. +66(0)22188357; Fax. +66(0)22188357  
E-mail address: [suchada.su@chula.ac.th](mailto:suchada.su@chula.ac.th)

### Introduction

The family Orchidaceae is one of the biggest flowering plant families. Many bioactive phytochemicals such as stilbenoids, flavonoids, terpenoids and alkaloids have been discovered from orchids.<sup>1-2</sup> *Paphiopedilum*, which is a genus of subfamily Cypripedioideae, consists of several lady's slipper orchids native to Thailand. Although these orchids are widely grown for their characteristic, beautiful flowers, their chemical constituents are very rarely investigated. This study aims to explore chemical constituents of the roots of *Paphiopedilum exul*, a terrestrial lady's slipper orchid endemic to southern Thailand.


### Material and methods

**General procedures:** <sup>1</sup>H-NMR (300 MHz), <sup>13</sup>C-NMR (75 MHz), and 2D-NMR spectra were obtained on a Bruker Avance DPX-300 NMR spectrometer (Bruker Corp., Billerica, MA, USA), in CDCl<sub>3</sub> or acetone-*d*<sub>6</sub> with tetramethylsilane

as an internal standard. Silica gel 60 F<sub>254</sub> plates (Merck, Darmstadt, Germany) were used for TLC analysis. Column chromatography (CC) was performed using either silica gel 60 (230-400 mesh, Merck) as the adsorbent or Sephadex LH-20 (Pharmacia Biotech AB, Uppsala, Sweden) as the gel filter.

**Plant material:** The whole plants of *Paphiopedilum exul* (Ridl.) Rolfe were purchased from Chatuchak market, Bangkok, in January 2015. Plant identification was performed by comparison with authentic specimen (QBG No. 13143) at the herbarium of the Botanical Garden Organization, Ministry of Natural Resources and Environment, Thailand. The fresh roots were separated from the plants, cleaned and dried at temperature not more than 50°C.

**Extraction and isolation:** Dried roots of *P. exul* (380 g) were ground and macerated three times with MeOH at room temperature. The MeOH extract was concentrated under reduced pressure and combined to give 100 g of crude MeOH extract. A portion of the extract (50 g) was separated on a silica gel column (1.25 kg, 10 × 40 cm), washed down with *n*-hexane/acetone (3:1), to yield eight fractions (A-H). Fraction C (2.9 g) was chromatographed on a silica gel column (150 g, 4.5 × 19 cm), eluted with CH<sub>2</sub>Cl<sub>2</sub>/acetone (40:1), to provide eight subfractions (C1-C8). Subfraction C2 (890 mg) was subjected to a Sephadex LH-20 CC using MeOH as the eluent to yield three subfractions (C21-C23). Purification of subfraction C23 (300 mg) by repeated silica gel CC, using CH<sub>2</sub>Cl<sub>2</sub>/acetone (40:1) as the eluent, afforded compound **1** (44 mg). Fraction E (9.9 g) was separated on a silica gel column (300 g, 4.5 × 40 cm) washed down with *n*-hexane/acetone (2:1) to give six subfractions (E1-E6). Subfraction E3 (3.4 g) was subjected to silica gel CC (175 g, 4.5 × 26 cm) eluted with CH<sub>2</sub>Cl<sub>2</sub>/acetone (30:1) to give nine subfractions (E31-E39). Compound **2** (32 mg) was obtained from subfraction E38. Subfraction E35 (1.4 g) was chromatographed on a silica gel column (70 g, 3 × 33 cm) eluted with CH<sub>2</sub>Cl<sub>2</sub>/acetone (30:1) to give three subfractions (E351-E353). Silica gel CC (55 g, 3 × 26 cm) of subfraction E352 (1.1 g), eluted with CH<sub>2</sub>Cl<sub>2</sub>/acetone (30:1), afforded five subfractions (E3521-E3525). A mixture of compounds **3** and **4** (72 mg) was obtained from subfraction E3525. Gel filtration of fraction E5 (780 mg) on a Sephadex LH-20 column eluted with MeOH yielded four subfractions (E51-E54). Subfraction E53 (290 mg) was chromatographed on a silica gel column (15 g, 2 × 14 cm), eluted with CH<sub>2</sub>Cl<sub>2</sub>/acetone (20:1), to give six subfractions (E531-E536). Compound **5** (58 mg) was obtained from subfraction E535.



**Figure 1.** Chemical structures of compounds **1-5**

## Results

Five *trans*-stilbenes were isolated from the MeOH extract of the roots of *Paphiopedilum exul*. They were identified as (*E*)-3-methoxy-5-hydroxystilbene (**1**), (*E*)-3,5'-dimethoxy-5,2'-dihydroxystilbene (**2**), thunalbene (**3**), (*E*)-3-methoxy-5,2'-dihydroxystilbene (**4**), and (*E*)-3-methoxy-2-(4-hydroxybenzyl)-5,2'-dihydroxystilbene (**5**), respectively, by comparison of their <sup>1</sup>H-NMR, <sup>13</sup>C-NMR (see **Table 1**) and 2D-NMR data with previous reports.<sup>3-5</sup>

## Discussion

The stilbenes isolated in this study represented interesting chemotaxonomic significance. Among these *trans*-stilbenes, (*E*)-3-methoxy-5-hydroxystilbene (**1**) is the most widely found one. It has been reported as a constituent of pteridophytes, gymnosperms, monocots and dicots,<sup>6-13</sup> and has been reported to display various biological activities<sup>(6). 14-17</sup> Its wide distribution might be due to its chemical structure, which is basic *trans*-stilbene with biosynthetically common oxy-substitutions at positions 3 and 5. On the other hand, thunalbene (**3**), with another hydroxy substitution at position 3', occurred more limitedly in nature and has been previously found in only two monocot families, i.e. Orchidaceae and Dioscoreaceae.<sup>5,18-19</sup> The

www.tjps.pharm.chula.ac.th

compound was also reported to be anti-inflammatory.<sup>20</sup> The structures of the other three isolated stilbenes from *P. exul*, i.e. (*E*)-3,5'-dimethoxy-5,2'-dihydroxystilbene (**2**), (*E*)-3-methoxy-5,2'-dihydroxystilbene (**4**), and (*E*)-3-methoxy-2-(4-hydroxybenzyl)-5,2'-dihydroxystilbene (**5**), similarly possess 2'-hydroxy substitution. They have once been isolated from a *Phragmipedium* orchid, which belongs to the same subfamily (Cypripedioideae) as *Paphiopedilum*.<sup>4</sup> This position of the hydroxy group is conveniently located for the formation of 2-phenylbenzofuran stilbenes found in the roots of another *Paphiopedilum* species, *P. godefroyae*.<sup>21</sup>

**Table 1.** <sup>13</sup>C-NMR (75 MHz) spectral data of compounds **1-5** (in CDCl<sub>3</sub>)

Position	1	2	3	4	5*
1	139.7	139.8	138.8	140.0	139.3
2	104.9	105.1	105.0	105.0	119.7
3	161.1	161.1	161.1	161.1	159.5
4	101.0	101.1	101.1	101.0	99.0
5	156.9	156.9	156.8	156.8	157.3
6	105.9	105.9	106.0	105.9	104.4
7	128.3	129.7	129.0	127.2	127.0
8	129.4	123.7	128.8	124.4	125.7
1'	137.0	125.1	139.5	123.7	125.3
2'	126.6	147.3	113.0	153.1	155.5
3'	128.7	116.9	155.8	116.0	120.4
4'	127.8	114.7	114.9	129.6	129.1
5'	128.7	153.9	129.9	121.1	116.4
6'	126.6	111.6	119.5	128.8	127.3
3-OCH <sub>3</sub>	55.4	55.4	55.4	55.4	55.6
5'-OCH <sub>3</sub>		55.4			
1''					30.0
2''					133.3
3''/7''					129.7
4''/6''					115.5
5''					155.8

\* in acetone-*d*<sub>6</sub>

## Conclusion

Five *trans*-stilbenes were obtained from the roots of a lady's slipper orchid, *Paphiopedilum exul*. The 2'-hydroxy substitution in three of these stilbenes appears to be characteristic of orchids in the subfamily Cypripodioideae.

## Acknowledgements

The authors would like to acknowledge partial financial support from a grant for International Research Integration: Research Pyramid from the Ratchadaphiseksomphot Endowment Fund.

## References

- Gutiérrez RMP. Orchids: A review of uses in traditional medicine, its phytochemistry and pharmacology. *J Med Plants Res.* 2010; 4: 592-638.
- Hossain MM. Therapeutic orchids: Traditional uses and recent advances — An overview. *Fitoterapia* 2011; 82: 102-40.
- Ngo K-S, Brown GD. Stilbenes, monoterpenes, diarylheptanoids, labdanes and chalcones from *Alpinia katsumadai*. *Phytochemistry* 1998; 47: 1117-23.
- Garo E, Hu J-F, Goering M, Hough G, O'Neil-Johnso M, Eldridge G. Stilbenes from the orchid *Phragmipedium* sp. *J Nat Prod.* 2007; 70: 968-73.
- Majumder P, Roychowdhury M, Chakraborty S. Thunalbene, a stilbene derivative from the orchid *Thunia alba*. *Phytochemistry* 1998; 49: 2375-8.
- Cao S, Radwan MM, Norris A, Miller JS, Ratovoson F, Mamisoa A, Andriantsiferana R, Rasamison VE, Rakotonandrasana S, Kingston DGI. Cytotoxic and other compounds from *Didymochlaena truncatula* from the

- Madagascar rain forest. J Nat Prod. 2006; 69: 284-6.
7. Suga T, Ohta S, Munesada K, Ide N, Kurokawa M, Shimizu M, Ohta E. Endogenous pine wood nematocidal substances in pines, *Pinus massoniana*, *P. strobus* and *P. palustris*. Phytochemistry 1993; 33: 1395-401.
  8. Fu Z, Zhang Y, Tan N. Chemical constituents of *Keteleeria evelyniana*. Nat Prod Res Dev. 2008; 20: 257.
  9. Starks CM, Williams RB, Norman VL, Lawrence JA, O'Neil-Johnson M, Eldridge GR. Phenylpropanoids from *Phragmipedium calurum* and their antiproliferative activity. Phytochemistry 2012; 82: 172-5.
  10. Hema P, Nair MS. Flavonoids and other constituents from the rhizomes of *Alpinia calcarata*. Biochem Syst Ecol. 2009; 37: 52-4.
  11. Qan-Kun Y, Bo L. New withanolides from *Nicandra physaloides* (Solanaceae). Plant Divers. 2012; 34: 101-6.
  12. Patel NK, Bhutani KK. Pinostrobin and cajanus lactone isolated from *Cajanus cajan* (L.) leaves inhibits TNF- $\alpha$  and IL-1 $\beta$  production: *in vitro* and *in vivo* experimentation. Phytomedicine 2014; 21: 946-53.
  13. Xu J, Tan N. Phenolic compounds from *Jatropha curcas*. Zhongguo Zhongyao Zazhi 2012; 37: 3074-7.
  14. Laavola M, Nieminen R, Leppänen T, Eckerman C, Holmbom B, Moilanen E. Pinosylvins and monomethylpinosylvins, constituents of an extract from the knot of *Pinus sylvestris*, reduce inflammatory gene expression and inflammatory responses *in vivo*. J Agric Food Chem. 2015; 63: 3445-53.
  15. Plumed-Ferrer C, Väkeväinen K, Komulainen H, Rautiainen M, Smeds A, Raitanen J-E, Eklund P, Willför S, Alakomi H-L, Saarela M, von Wright A. The antimicrobial effects of wood-associated polyphenols on food pathogens and spoilage organisms. Int J Food Microbiol. 2013; 164: 99-107.
  16. Araujo MFd, Vieira IJC, Braz-Filho R, Vieira-da-Motta O, Mathias L. Chemical constituents from *Swartzia apetala* Raddi var. *glabra* and evaluation of their antifungal activity against *Candida* spp. Rev Bras Farmacogn. 2009; 19: 366-9.
  17. Sobolev VS, Khan SI, Tabanca N, Wedge DE, Manly SP, Cutler SJ, Coy MR, Becnel JJ, Neff SA, Gloer JB. Biological activity of peanut (*Arachis hypogaea*) phytoalexins and selected natural and synthetic stilbenoids. J Agric Food Chem. 2011; 59: 1673-82.
  18. Yuan Z-P, Zhong Y-J, Su X-H, Li C-Y, Du H-J, Wen Y-Y, Li Y-F, Liang B. Chemical constituents of *Pholidota leveilleana*. Biochem Syst Ecol. 2013; 49: 148-51.
  19. Liu H, Tsim KW, Chou GX, Wang JM, Ji LL, Wang ZT. Phenolic compounds from the rhizomes of *Dioscorea bulbifera*. Chem Biodivers. 2011; 8: 2110-6.
  20. Wang J, Matsuzaki K, Kitanaka S. Stilbene derivatives from *Pholidota chinensis* and their anti-inflammatory activity. Chem Pharm Bull. 2006; 54: 1216-8.
  21. Lertnitikul N, Jittham P, Khankhampoch L, Pattamadilok C, Sukrong S, Suttisri R. Cytotoxic stilbenes from the roots of *Paphiopedilum godefroyae*. J Asian Nat Prod Res. (submitted).