



# Synthesis, characterization, and anti-tumor application of a novel Zinc(II)-L-ascorbic acid derivative

Mohammed A. Mohammed<sup>1</sup>, Mouhaned Y. Al-Darwesh<sup>2</sup>,  
Rasim Farraj Muslim<sup>3</sup>, Muwafaq Ayesh Rabeea<sup>4\*</sup>

<sup>1</sup>Department of Ramadi Education, Directorate of Education in Anbar, Ministry of Education, Ramadi Anbar, Iraq, <sup>2</sup>Department of Chemistry, College of Science, University of Anbar, Anbar, Iraq, <sup>3</sup>Department of Environmental, College of Applied Sciences-Hit, University of Anbar, Anbar, Iraq, <sup>4</sup>Department of Applied Chemistry, College of Applied Sciences-Hit, University of Anbar, Anbar, Iraq

## Corresponding Author:

Muwafaq Ayesh Rabeea,  
Department of Applied  
Chemistry, College of Applied  
Sciences-Hit, University of  
Anbar, Anbar, Iraq. E-mail:  
muw88@uoanbar.edu.iq

**Received:** March 10, 2021

**Accepted:** October 08, 2021

**Published:** March 23, 2022

## ABSTRACT

In this research, a derivative of L-ascorbic acid (L) as a ligand was synthesized from the reaction of this acid with acetone compound in an acidic medium, and then, a complex reaction was carried out between the synthesized derivatives with zinc chloride salt using ethanol as a solvent. This is a novel attempt to use 5,6-O-isopropylidene-L-ascorbic acid as a derivative of L-ascorbic acid in the synthesis of zinc complex  $[Zn(L)Cl_2]$ . Thin-layer chromatography was used to determine the time required for the completion of the reactions. The formation of the derivative (L) was confirmed by m.p, Rf, Fourier transform infrared (FT-IR), <sup>1</sup>H-nuclear magnetic resonance, and mass spectra, and the synthesis of  $[Zn(L)Cl_2]$  was proved by m.p, Rf, FT-IR, and mass spectra. The biological activity of the fabricated complex was evaluated against three human cancer cell lines, including prostate cancer cell line (PC-3), colon cancer cell line (Caco-2), and breast cancer cell line (MCF-7). The  $[Zn(L)Cl_2]$  exhibited a good reduction in cell viability in a dose-dependent system, where cell viability is reduced by increasing the concentration of the complex. In addition, its antioxidant activity was investigated and demonstrated significant activity at concentrations of 100 and 200 g/mL. Hence, this complex has the potential to be a promising drug for the treatment of PC-3, Caco-2, and MCF-7.

**Keywords:** Antioxidant, Ascorbic acid, Breast cancer, Zn(II) complex

## INTRODUCTION

The Vitamin C discovery and its medical application go back to the 16<sup>th</sup> and 17<sup>th</sup> centuries. It was used to prevent the “scurvy” disease among sailors in Europe. Recently, the variety of Vitamin C is used in many fields such as pharmaceuticals, industry, food, and agriculture.<sup>[1]</sup> For many physiological functions, Vitamin C is required by the human body, and the functions of this vitamin are most important and represented by enhancement of response in body immune system, iron absorption, and pulmonary function.<sup>[2]</sup> Extensive clinical, animal, and epidemiological studies were conducted to examine the role of ascorbic acid for the prevention of different types of cancers. Except for gastric cancer, the results were found to be incomplete.<sup>[3]</sup>

Despite the fact of the simple structure of this molecule, the biochemistry of Vitamin C is not fully understood because

of its complicated redox chemistry. For this reason, this molecule mostly is used in inorganic chemistry as a reducing agent. Udenfried *et al.*<sup>[4]</sup> have studied the interaction between metal ions and this vitamin almost 50 years ago. This work was about Vitamin C oxidation by oxygen gas (O<sub>2</sub>). Other studies have been carried out on the interactive nature of ascorbic acid and metal ions. Martell *et al.*<sup>[5]</sup> examined the catalytic role of metals in the oxidation of L-ascorbic. The zinc-Vitamin C interaction is also very important, as Zn(II) is necessary for the activity of over 200 metallic enzymes in the human body.<sup>[6]</sup>

Zinc is one of the outstanding trace elements that have a crucial role in different cellular processes including cell proliferation, free radicals defense, and differentiation.<sup>[7,8]</sup> Zinc is an essential element in the structure of many enzymes and proteins, including cellular signaling proteins, transcription factors, and repair enzymes of DNA.<sup>[9,10]</sup> According to the

literature, zinc ion has been shown to have an important role in the regulation of growth in mammalian cells.<sup>[11-13]</sup> Constantly, the lowered Zn levels are associated with systemic abnormalities including cancer development.<sup>[14]</sup> Therefore, zinc complexes have been appeared with their distinctive properties as potential anticancer agents.<sup>[15]</sup>

However, this work aims to use one of the derivatives of L-ascorbic acid (L) as a vital ligand for the fabrication of [Zn(L)Cl<sub>2</sub>]. The theoretical and spectroscopic analyses indicate the formation of the target complex. The antioxidant activity of the synthesized complex was evaluated, as well as its anti-tumor activity against prostate cancer cell line (PC-3), colon cancer cell line (Caco-2), and breast cancer cell line (MCF-7).

## EXPERIMENTAL

### Synthesis of 5,6-O-isopropylidene-L-ascorbic Acid

A saturated solution of L-ascorbic acid derivative (10.00 g, 57.00 mmol) in 100 ml of a mixture of absolute acetone and hydrochloric acid (V/V) was stirred at room temperature for 20 min. After that, 80 ml of n-hexane was added to the solution. The residue was washed 4 times with acetone-hexane (4:7 V/V); then, the solvent was removed under reduced pressure to yield a white crystallized L-ascorbic acid derivative.<sup>[16]</sup>

### Preparation of Complex [Zn(L)Cl<sub>2</sub>]

A 15 mL of L-ascorbic acid (0.2 g, 0.92 mmol) dissolved in ethanol were added to 15 mL of [ZnCl<sub>2</sub>] (0.125 g, 0.92 mmol) suspended in ethanol. The mixture was refluxed for 2 h, and the formed brown precipitate was filtered and dried at room temperature.<sup>[17,18]</sup>

### Characteristics of the L-ascorbic Acid Derivative

L-ascorbic acid derivative (C<sub>9</sub>H<sub>12</sub>O<sub>6</sub>): Color (White). Yield (10.96 g, 91.33 %). m. p. (218–221°C). R<sub>f</sub> (0.61). Fourier transform infrared (FTIR) (cm<sup>-1</sup>): Broad band at 2400–3600 for carboxylic acid hydroxyl (O-H); 2995 ν(C-H); 1756 ν(C=O); 1664 ν(C=C); 1141 w(C-O). <sup>1</sup>H-nuclear magnetic resonance (NMR) (400 MHz, d<sub>6</sub>-DMSO): (ppm): 11.27 (s, for the proton of carboxylic acid hydroxyl OH [G group]), 8.46 (s, for alcohol hydroxyl OH [F group]), 4.71 (s, for the proton of OH [E group]), 4.25 and 4.26 (d, for the proton of CH [D group]), 4.07, 4.09, 4.11, and 4.12 (q, for the proton of the CH [C group]), 3.86 and 3.88 (d, for the 2 proton of the CH<sub>2</sub> [B group]), 1.26 (s, for 6 protons of 2CH<sub>3</sub> [A groups]). Mass (m/z): 216.0 (C<sub>9</sub>H<sub>12</sub>O<sub>6</sub>), 159.0 (C<sub>7</sub>H<sub>12</sub>O<sub>4</sub><sup>+</sup>), 130.0 (C<sub>6</sub>H<sub>10</sub>O<sub>3</sub><sup>+</sup>), 88 (C<sub>4</sub>H<sub>8</sub>O<sub>2</sub><sup>+</sup>), 57.1 (C<sub>2</sub>H<sub>2</sub>O<sub>2</sub><sup>+</sup>), 42.0 (C<sub>3</sub>H<sub>6</sub><sup>+</sup>), 30.1 (CH<sub>2</sub>O<sup>+</sup>). Microanalysis (The microanalysis of elements for prepared complex was estimated by Shimadzu-AA-670-Flame spectrophotometer in the laboratories of Gazi University in Turkey): 50.0 (C Theoretical), 50.6 (C Practically), 5.60 (H Theoretical), and 5.88 (H Practically).

### Characteristics of the Prepared Complex

Perpetrated complex ([Zn(L)(Cl)<sub>2</sub>]): Color (Brown). Yield (0.28 g, 86.41%). m. p. (288–289°C), R<sub>f</sub> (0.56). FTIR (cm<sup>-1</sup>):

2981, 2894 (C-H); 1743 (C=O); 1612 (C=C); 1139 (C-O); 2400–3610 (O-H). Mass (m/z): 350.2 (C<sub>9</sub>H<sub>11</sub>Cl<sub>2</sub>O<sub>6</sub>Zn<sup>+</sup>), 289.8 (C<sub>4</sub>H<sub>5</sub>Cl<sub>3</sub>O<sub>4</sub>Zn<sup>+</sup>), 156.0 (C<sub>7</sub>H<sub>10</sub>ClO<sub>2</sub><sup>+</sup>), 133.8 (C<sub>4</sub>H<sub>3</sub>ClO<sub>3</sub><sup>+</sup>), 84.0 (C<sub>4</sub>H<sub>6</sub>O<sub>2</sub><sup>+</sup>), 71.9 (C<sub>3</sub>H<sub>4</sub>O<sub>2</sub><sup>+</sup>), 58.1 (C<sub>3</sub>H<sub>6</sub>O<sup>+</sup>). Microanalysis: 30.58 (C Theoretical), 29.9 (C Practically), 2.88 (H Theoretical), 2.11 (H Practically), 20.23 (Cl Theoretical), 19.52 (Cl Practically), 18.66 (Zn Theoretical), and 17.92 (Zn Practically).

## Biological Activity

Antioxidant activity of the prepared complex was detected using 2,2-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging assay according to Rajesh and Natvar.<sup>[19]</sup>

The cytotoxic effect of the complex was conducted using ready-to-use MTT kit.<sup>[20]</sup> The kit contents are: (1 mL × 10 vials MTT solution) and (50 mL × 2 bottle solubilizing solution).

## Statistical Analysis

Statistical Analysis System (2012) program was employed for examining the different factors in the study parameters. To compare the means and Chi-square, test least significant difference–LSD test analysis of variance was used to compare between percentages. The estimation of correlation coefficient between variables has also been studied.<sup>[21]</sup>

## RESULTS AND DISCUSSION


### L-ascorbic Acid Derivative

L-ascorbic acid molecule has four (-O-H) groups, all these groups are active for known esterification reactions and other reactions. L-ascorbic acid derivatives synthesis at 2- and 3-position requires the conversion of L-ascorbic acid into its 5,6-isopropylidene derivative because the carbon-6 hydroxyl group (a primary hydroxyl group) is the most reactive group.<sup>[22]</sup> The acetal group is readily hydrolyzed by a dilute acid.<sup>[23,24]</sup> L-ascorbic derivative was synthesized from the reaction of acetone and L-ascorbic acid in acidic medium, according to the literature,<sup>[16]</sup> Scheme 1.

In the first step, the suggested mechanism of the synthesized derivative involved protonation, the second step was to attack the electron pair of the hydroxyl oxygen in the terminal continuous chain on the carbonyl carbon of the ketone, and in the third step, the transfer of a proton from the terminal hydroxyl group to the nearest hydroxyl group, followed by a fourth step by losing a water molecule and cyclic closure and in the fifth step occurred opening of the ring containing lactone group (O-C=O), Scheme 2:

The FTIR spectrum of L-ascorbic acid derivative [Figure 1] showed a broad band at 2400–3600 cm<sup>-1</sup> for (O-H) of carboxylic acid and other bands were observed at 2995 cm<sup>-1</sup> for (C-H aliphatic), 1756 cm<sup>-1</sup> for (C=O) of carboxylic acid, 1664 cm<sup>-1</sup> for aromatic (C=C), and 1141 cm<sup>-1</sup> for (C-O).<sup>[25,26]</sup>

The <sup>1</sup>H-NMR spectrum of an L-ascorbic acid derivative [Figure 2] reveals a singlet signal at 11.27 ppm that corresponds to the proton of the hydroxyl group OH of carboxylic acid (g group), a singlet signal at 8.46 ppm for the proton of OH (f group), a singlet signal at 4.71 ppm for the proton of OH




### The Synthesis of Complex $[\text{Zn}(\text{L})(\text{Cl})_2]$

(86.41%) of  $[\text{Zn}(\text{Cl})_2]$  was obtained with a brown color and a melting point of 288–289°C, Scheme 3.

The FTIR spectrum of  $[\text{Zn}(\text{L})(\text{Cl})_2]$  complex shows stretching band at  $2981\text{ cm}^{-1}$  asymmetric,  $2894\text{ cm}^{-1}$  symmetric for (C-H aliphatic),  $1743\text{ cm}^{-1}$  for (C=O) of carboxylic acid,  $1612\text{ cm}^{-1}$  for (C=C),  $1139\text{ cm}^{-1}$  for (C-O), and broad band at  $2400\text{--}3610\text{ cm}^{-1}$  for (O-H) of carboxylic acid,<sup>[27,28]</sup> Figure 4 of FTIR spectra of  $[\text{Zn}(\text{Asc})(\text{Cl})_2]$  complex.

The mass spectrum of the complex [Figure 5] shows the following signals in Table 2. These signals are representing the molecular weight of the complex in addition to the emergence of multiple fissures (structure of  $[Zn(L)Cl_2]$  complex and its fragmentation), which confirm the formation of this complex with the proposed formulas and the suggested molecular weights that are consistent with the other results.

A microanalysis of the synthesized L-ascorbic acid derivative and  $[\text{Zn}(\text{L})\text{Cl}_2]$  complex elements was measured and a match was found between theoretical percentages and the calculated practical percentages.



**Figure 3:** The mass spectrum of L-ascorbic acid derivative


**Table 1:** Mass spectra and fragmentation of L-ascorbic acid derivative

Molecule	Structure of fragments	Fragmentation chemical formula	Mass/charge: m/z
		$C_9H_{13}O_6^{+}$	216.0
		$C_7H_{10}ClO_2^{+}$	159.0
		$C_4H_3ClO_3^{+}$	130.0
		$C_4H_8O_2^{+}$	88.0
		$C_2H_2O_2^{+}$	57.1
		$C_3H_6^{+}$	42.0
		$CH_2O^{+}$	30.1

## Quantum Chemical Properties

The global chemical reactivity indices of  $[Zn(L)Cl_2]$  were revealed by molecular orbital calculations depending on lowest unoccupied molecular orbital (LUMO) and highest occupied molecular orbital (HOMO) calculations [Table 3]. These agents are useful for predicting the interaction of  $[Zn(L)Cl_2]$  with biological molecules in a human body and their diffusion across cell membranes and the brain-blood barrier.<sup>[29,30]</sup> Figure 6 shows that the energy difference between

HOMO and LUMO of  $[Zn(L)Cl_2]$  is 4.400 eV, so there is a contribution of the prepared complex on the ability to accept electrons. Related parameters such as electronic chemical potential ( $\mu$ ), chemical hardness ( $\eta$ ), and electrophilicity index ( $\omega$ ) were estimated using Equations 1, 2, and 3, which found to be 2200 eV, 3930 eV, and 1786 eV, respectively. These values reveal the biologically pharmacodynamic activity of  $[Zn(L)Cl_2]$ . Furthermore, the negative value (-3.459) of the solubility class (Log S) means having the ability to dissolve rapidly with serum and proteins. Finally, using Swiss ADME




**Figure 4:** Fourier transform infrared of  $[Zn(L)(Cl)_2]$  complex


**Table 2:** Mass/Charge (m/z) of the fragmentation of  $[Zn(L)(Cl)_2]$  complex

$[Zn(Asc)(Cl)_2]$ complex	Structure of fragments	Chemical formula of complex fragments	Mass/charge: m/z
		$C_9H_{11}Cl_2O_6Zn^{+}$	350.2
		$C_4H_5Cl_3O_4Zn^{++}$	289.8
		$C_7H_{10}ClO_2^{+}$	156.0
		$C_4H_3ClO_3^{+}$	133.8
		$C_4H_6O_2^{+}$	84.0
		$C_3H_4O_2^{+}$	71.9
		$C_3H_6O^{+}$	58.1





**Figure 5:** Mass spectrum of  $[Zn(L)(Cl)_2]$  complex



**Figure 6:** (a and b) Frontier molecular orbitals of  $[Zn(L)Cl_2]$

online, the prepared complex was effective on drug-likeness criteria derived from the rules of Lipinski, Ghosh, and Veber.<sup>[31]</sup>

$$\mu = (\varepsilon_{HOMO} + \varepsilon_{LUMO}) / 2 \quad (1)$$

$$\eta = (\varepsilon_{LUMO} - \varepsilon_{HOMO}) / 2 \quad (2)$$

$$\omega = \mu^2 / 2\eta \quad (3)$$

### Antioxidant Activity of $[Zn(L)Cl_2]$ Complex

The activity of free radical scavenging of the synthesized complex has been studied using the stable (DPPH) of free radical scavenging assay. DPPH is characterized as a stable

**Table 3:** Physicochemical properties of  $[Zn(L)Cl_2]$

$[Zn(L)Cl_2]$	Physicochemical properties
-6.130	HOMO (eV)
-1.730	LUMO (eV)
	Dipole moment (debye)
-3.459	Log S
2.982	Log P
4.400	Energy gap ( $\Delta$ ) (eV)
3.930	$\mu$ (eV)
2.200	$\eta$ (eV)
1.786	$\omega$ (eV)



radical due to the delocalization of the maximum absorption about 517 nm and the spare electron. The activity of DPPH in scavenging has been commonly used to estimate the antiradical activities of different compounds,<sup>[32-34]</sup> Table 4.

Results in Table 4 revealed that 200  $\mu\text{g/mL}$  of complex had the highest scavenging activity (86.03%) that is showing the more concentration of the complex, and the more free radicals scavenging percentage. When compared to free radical scavenging positive antioxidant ascorbic acid activity, the prepared complex exhibits nearly the same pattern of free radical scavenging activity, particularly at concentrations of 12.5–50  $\text{g/mL}$  (Figure 7). While the concentrations of 100 and 200  $\mu\text{g/mL}$  showed higher activities compared with ascorbic acid (as a positive control). Free radicals play a critical role developing some health conditions, example cataracts, cancer, and cardiovascular disease. Free radicals have been represented in both acceleration and initiation of the aging process.<sup>[35]</sup> It is well-known that free radicals are the main cause of oxidation and that they can be prevented or controlled by the use of various antioxidants.<sup>[36]</sup> Many chemical compounds possess

**Table 4:** Antioxidant activity of [Zn (L) Cl<sub>2</sub>] complex using 2,2-Diphenyl-2-Picrylhydrazyl method

Compound	Compound				
	12.5 (µg/mL)	25 (µg/mL)	50 (µg/mL)	100 (µg/mL)	200 (µg/mL)
Ascorbic acid	17.63±7.196	40.43±7.081	54.48±2.412	61.77±1.351	72.11±3.002
Complex	22.90±1.836 (NS)	39.00±1.732 (NS)	57.60±2.207 (NS)	74.07±1.007**	86.03±4.028**

\*\*Significant at level ( $P < 0.001$ ). NS: Non-significant


**Scheme 1:** Synthesis of L-ascorbic acid derivative

**Scheme 2:** Suggested mechanism of L-ascorbic acid derivative preparation

great antioxidant activities with capabilities of reducing the oxidative reactions in cells due to their ability to donate

hydrogen atoms, therefore, they are considered useful in the treatment of many diseases including inflammatory, cancer,

**Table 5:** Anticancer activity of [Zn (L) Cl<sub>2</sub>] complex on MCF-7, prostate cancer-3, and Caco-2 cell lines after 24 h of incubation at 37°C using the MTT method


	Concentration (µg/mL)	400	200	100	50	25	12.5	6.2	IC <sub>50</sub>	P
MCF7		44.445±1.208	47.801±1.947	51.235±2.032	61.458±1.365	75.502±2.453	85.764±1.445	94.522±0.571	72.46	0.0001>
WRL68		78.665±2.549	81.636±1.438	86.150±1.275	90.394±1.029	93.904±1.112	94.985±0.678	94.753±0.964	248.5	
PC3		61.574±3.478	65.548±1.237	70.447±1.993	79.012±2.166	86.883±1.563	94.252±1.142	94.830±0.241	128.2	<0.0001
WRL68		73.349±1.474	81.991±5.005	92.477±1.505	93.866±0.613	95.062±0.177	94.753±0.571	95.216±0.241	470.1	
Caco-2		32.793±7.442	37.847±1.736	63.812±2.744	90.046±2.425	96.219±0.678	96.644±1.365	97.26±0.3	98.70	<0.0001
WRL68		70.409±1.414	84.028±1.002	91.667±1.621	95.332±1.183	95.216±0.821	95.949±1.028	95.45±0.3	333.7	

PC: Prostate cancer, Caco: Colon cancer cell line (Caco-2), MCF: Breast cancer cell line


and cardiovascular diseases.<sup>[37]</sup> Zn complexes are more active DPPH scavengers than the corresponding free compounds.<sup>[38]</sup> A zinc complex of 2-hydroxy-salicylhydrazide-isatin hydrazone has two free hydroxyl groups in its [ZnL<sub>2</sub>] formula which exhibited good radical scavenging activity (SC<sub>50</sub> = 31.13 µM). Özdemir<sup>[39]</sup> suggested that this higher activity might be dependent on the stabilization of radical with the phenolic moieties of zinc complex. In addition, Andelescu *et al.* reported that Zn(II) complexes containing four free hydroxyl groups in their formulae which showed remarkable scavenging activity (SC<sub>50</sub> = 8.10–12.17 µM).<sup>[40]</sup>

## Effect of [Zn(L)Cl<sub>2</sub>] Complex on Different Cancer Cell Lines Using the MTT Method

3-(Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay was achieved to evaluate the cytotoxic effect of the complex against three different types of cancer cell lines; Caco-2, PC-3, and MCF-7. This assay was conducted to determine the cell viability and the rate of inhibition by the implementation of different concentrations of the complex on the targeted cancer cell lines. For the most significant IC<sub>50</sub>




**Figure 7:** Antioxidant activity of [Zn(l)(Cl)<sub>2</sub>] complex using DPPH method



**Figure 8:** Cytotoxicity effect of [Zn(l)(Cl)<sub>2</sub>] complex on breast cancer cell line-7 (a), prostate cancer-3 (b), and colon cancer cell line-2 (c) cell lines after 24 h of incubation at 37°C





**Scheme 3:** The synthesis of  $[Zn(L)(Cl)_2]$  complex

values, the best values were chosen. MTT colorimetric assay was used to determine the cell viability at every time point. The cytotoxic effect of the complex toward three different types of cancer cell lines is reported in Table 5 and Figure 8.

Results indicated that the incubation of tumor cell lines with the complex at different concentrations ranged between 12.5 and 400  $\mu\text{g/mL}$  for 24 h showed that viability of cell reduced in a dose-dependent pattern by the increase of the complex concentration. The cell viability percentage (%) of a lowest value was recorded at the concentration of 400  $\mu\text{g/mL}$ . The explanation of this case is still unrevealed, but it is known that significant amounts of the complex are hydrolyzed outside the cell and excreted by renal on plasma proteins binding, with only few DNA binds. It may be attributed to the strong Zn-O bond, which may result in more cell penetration by the complex and becoming bound to DNA. Significant differences were observed in the complex inhibition rate among different cell lines by the calculating of the  $IC_{50}$ . The complex showed the most cytotoxic activity on the MCF-7 cells with an  $IC_{50}$  value of 72.46  $\mu\text{g/mL}$ , whereas  $IC_{50}$  values of PC-3 and Caco-2 were 128.2 and 98.70  $\mu\text{g/mL}$ , respectively.

Researchers have paid attention toward designing metal complexes as anticancer agents. The metal complexes designing process depends on the donation ability and the metal ion type along with its oxidation state.<sup>[41]</sup> Cancer cell toxicity varied from one cell line to another, due to different types of cancer cell lines in their receptors the presence of the complex affected certain receptors on the surfaces of these cells, and through these receptors, the cells are responsive to the action of many substances that stimulate apoptosis.<sup>[42]</sup> Particular zinc thiosemicarbazone complexes have been studied and confirmed to have anti-cancer activity as antioxidant agents. They have shown cytotoxicity similar to cisplatin and significantly effective against cancer cell lines. The zinc complexes seem to be concentrated in a relatively small zone of the cell with a low general uptake in the rest of the cytoplasm. However, the cell absorption of mineral complexes depends on the cell type.<sup>[43]</sup>

## CONCLUSION

The formation of both the L-ascorbic acid derivative and  $[Zn(L)(Cl)_2]$  was demonstrated by spectroscopic studies (FTIR,  $^1\text{H-NMR}$ , and mass spectrum). This structure had a significant effect on the cancer cells under evaluation. DPPH is characterized as a stable radical due to the delocalization of the maximum absorption about 517 nm and the spare electron. Significant differences were observed in the complex inhibition rate among different cell lines by the calculating of the  $IC_{50}$ . The

complex showed the most cytotoxic activity on the MCF-7 cells with a  $IC_{50}$  value of 72.46  $\mu\text{g/mL}$ , whereas  $IC_{50}$  values of PC-3 and Caco-2 were 128.2 and 98.70  $\mu\text{g/mL}$ , respectively.

## ACKNOWLEDGMENTS

Special thanks to president of the University of Anbar Prof. Dr. Mushtak Talib Al-Ouqaili. Thankful to the staff of environmental department in College of Applied Sciences-Hir and the staff of chemistry department in College of Sciences in University of Anbar. Thanks to Assist. Prof. Dr. Mustafa Nadhim Owaid from the Department of Hit Education, Directorate of Education in Anbar, Ministry of Education.

## REFERENCES

- Seib PA, Tolbert BM, editors. *Ascorbic Acid: Chemistry, Metabolism, and Uses*. Washington, USA: American Chemical Society; 1982.
- Weber P, Bendich A, Schalch W. La vitamina C y la salud humana, una revisión de los datos recientes correspondientes a las necesidades humanas. *Int J Vitam Nutr Res* 1996;66:19-30.
- Story JA, Kritchevsky D, Carroll KK. Kenneth K. Carroll (1923-1998). *J Nutr* 2000;130:1089-90.
- Udenfriend S, Clark CT, Axelrod J, Brodie BB. Ascorbic acid in aromatic hydroxylation. *J Biol Chem* 1954;208:731-9.
- Du J, Cullen JJ, Buettner GR. Ascorbic acid: Chemistry, biology and the treatment of cancer. *Biochim Biophys Acta* 2012;1826:443-57.
- Tajmir-Riahi HA. Coordination chemistry of Vitamin C. Part II. Interaction of L-ascorbic acid with  $Zn(II)$ ,  $Cd(II)$ ,  $Hg(II)$ , and  $Mn(II)$  ions in the solid state and in aqueous solution. *J Inorg Biochem* 1991;42:47-55.
- Ho E. Zinc deficiency, DNA damage and cancer risk. *J Nutr Biochem* 2004;15:572-8.
- Stefanidou M, Maravelias C, Dona A, Spiliopoulos C. Zinc: A multipurpose trace element. *Arch Toxicol* 2006;80:1.
- Prasad AS. Zinc: An overview. *Nutrition* 1995;11 1 Suppl:93-9.
- Prasad AS, Kucuk O. Zinc in cancer prevention. *Cancer Metastasis Rev* 2002;21:291-5.
- Fraker PJ, Lill-Elghanian DA. The many roles of apoptosis in immunity as modified by aging and nutritional status. *J Nutr Health Aging* 2004;8:56-63.
- Chang KL, Hung TC, Hsieh BS, Chen YH, Chen TF, Cheng HL. Zinc at pharmacologic concentrations affects cytokine expression and induces apoptosis of human peripheral blood mononuclear cells. *Nutrition* 2006;22:465-74.
- Franklin RB, Costello LC. The important role of the apoptotic effects of zinc in the development of cancers. *J Cell Biochem* 2009;106:750-7.
- Murakami M, Hirano T. Intracellular zinc homeostasis and zinc signaling. *Cancer Sci* 2008;99:1515-22.
- Kraljević TG, Wittner K, Lovrić I, Makuc D. Synthesis, structural studies, and cytostatic evaluation of 2,6-di-O-modified L-ascorbic acid derivatives. *Carbohydr Res* 2006;341:433-42.
- Shi W, Y. Chaturvedi KK, Kresge AJ, Yin Y. Acid-catalyzed hydrolysis of vinyl acetals. Reaction through the acetal rather than the vinyl ether functional group. *J Am Chem Soc* 1989;111:7185-90.
- Fife TH, Jao LK. General acid catalysis of ascorbic acid hydrolysis. *J Am Chem Soc* 1968;90:4081-5.
- Harfini ST, Muslim RF, Awad MA. Characterization and biological effectiveness of synthesized complexes of Palladium (II) from imine compounds. *Egypt J Chem* 2022;65:385-96.
- Davis J, Dilworth JR, Donnelly JS, Babson PI, Nithinvala A, et al. Fluorescence resonance energy transfer distribution of zinc bis(thiosemicarbazone) complexes in human cancer cells. *Chem Commun* 2005;7:845-7.
- Goparakrishnan SB, Kumarasi T, Subramanian IC. Comparative DFT study of phytochemical constituents of the fruits of *Cucumis trigonus* Roxb. and *Cucumis sativus* Linn. *J Comput Methods Phys Sci* 2013;2013:1-5.

15. Frezza M, Hindo S, Chen D, Davenport A, Schmitt S, Tomco D, et al. Novel metals and metal complexes as platforms for cancer therapy. *Curr Pharm Des* 2010;16:1813-25.
16. Al-Dolaymi MA, Al-Rawi KF, Al-Shamary FT, Al-Halbosi MM. Complex synthesis of Cis-[Pt (Asc)(NH<sub>3</sub>)<sub>3</sub>] and its effect on human breast cancer MCF-7 cell *in vitro*. *Indones J Chem* 2019;19:245-53.
17. Mahmood RS, Awad MA, Muslim RF. Preparation of some metallic complexes from selected organic ligands and characterization it spectrophotometry as a model of a ligand scavenger to metals in health applications. *Ann Trop Med Public Health* 2020;23:231374.
18. Ibrahim NS, Awad MA, Muslim RF. Synthesis and characterization of novel cadmium complexes from organic ligands. *Biochem Cell Arch* 2020;20:4079-89.
19. Patel Rajesh M, Patel Natvar J. *In vitro* antioxidant activity of coumarin compounds by DPPH, Super oxide and nitric oxide free radical scavenging methods. *J Adv Pharm Educ Res* 2011;1:52-68.
20. Freshney RI. *Culture of Animal Cells: A Manual of Basic Technique and Specialized Applications*. Hoboken, NJ: John Wiley & Sons; 2015.
21. Delwiche LD, Slaughter SJ. *The Little SAS Book: A Primer*. Cary, NC: SAS Institute; 2019.
22. Kraljević TG, Wittine K, Lovrić I, Makuc D. Synthesis, structural studies, and cytostatic evaluation of 5,6-di-O-modified L-ascorbic acid derivatives. *Carbohydr Res*. 2006;341:433-42.
23. Chiang Y, Chwang WK, Kresge AJ, Yin Y. Acid-catalyzed hydrolysis of vinyl acetals. Reaction through the acetal rather than the vinyl ether functional group. *J Am Chem Soc* 1989;111:7185-90.
24. Fife TH, Jao LK. General acid catalysis of acetal hydrolysis. The hydrolysis of 2-aryloxytetrahydropyrans. *J Am Chem Soc* 1968;90:4081-5.
25. Ayfan AK, Muslim RF, Noori NS. Preparation and characterization of novel disubstituted 1,3-oxazepine-tetraone from Schiff bases reaction with 3-methylfuran-2,5-dione and 3-phenyldihydrofuran-2,5-dione. *Res J Pharm Tech* 2019;12:1008-16.
26. Muslim RF, Saleh MM, Saleh SE. Synthesis and characterization of new sulphur six-membered heterocyclic compounds and evaluation their biological activity. *Rev AUS* 2019;26:129-35.
27. Hafidh SH, Muslim RF, Awad MA. Characterization and biological effectiveness of synthesized complexes of Palladium (II) from imine compounds. *Egypt J Chem* 2022;65:385-96.
28. Silverstein RM, Bassler GC. Spectrometric identification of organic compounds. *J Chem Educ* 1962;39:546.
29. Cherappanath RJ, John VD. Synthesis, characterization, anti-corrosive/anti-tumor applications of a new type arylazo compound 1, 7-bis (2-hydroxy phenyl)-4-(phenyl hydrazono)-hepta-1, 6-diene-3, 5-dione. *J Bio Tribocorros* 2018;4:1-13.
30. Gopalakrishnan SB, Kalaiarasi T, Subramanian R. Comparative DFT study of phytochemical constituents of the fruits of *Cucumis trigonus* Roxb. and *Cucumis sativus* Linn. *J Comput Methods Phys* 2014;2014:623235.
31. Abdullah JA, Aldahham BJ, Rabeea MA, Asmary FA, Alhajri HM, Islam MA. Synthesis, characterization and *in-silico* assessment of novel thiazolidinone derivatives for cyclin-dependent kinases-2 inhibitors. *J Mol Struct* 2021;1223:129311.
32. Hou JP, Wu H, Wang Y, Weng XC. Isolation of some compounds from nutmeg and their antioxidant activities. *Czech J Food Sci* 2012;30:164-70.
33. Muslim RF, Owaid MN. Synthesis, characterization and evaluation of the anti-cancer activity of silver nanoparticles by natural organic compounds extracted from *Cyperus sp.* rhizomes. *Acta Pharm Sci* 2019;57:126-46.
34. Owaid MN, Naeem GA, Muslim RF, Oleiwi RS. Synthesis, characterization and antitumor efficacy of silver nanoparticle from *Agaricus bisporus* Pileus, Basidiomycota. *Walailak J Sci Technol* 2020;17:75-87.
35. Simon JE, Wang M, Gbewonyo K, Rafi MM, Acquaye DF, Asianowa Y. U.S. Patent No. 7,371,413. Washington, DC: U.S. Patent and Trademark Office; 2008.
36. Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, Gluud C. Mortality in randomized trials of antioxidant supplements for primary and secondary prevention: Systematic review and meta-analysis. *JAMA* 2007;297:842-57.
37. Krishnaiah D, Sarbatly R, Nithyanandam R. A review of the antioxidant potential of medicinal plant species. *Food Bioprod Process* 2011;89:217-33.
38. Psomas G. Copper (II) and zinc (II) coordination compounds of non-steroidal anti-inflammatory drugs: Structural features and antioxidant activity. *Coord Chem Rev* 2020;412:213259.
39. Özdemir Ö. Bis-azo-linkage Schiff bases-Part (II): Synthesis, characterization, photoluminescence and DPPH radical scavenging properties of their novel luminescent mononuclear Zn (II) complexes. *J Photochem Photobiol A Chem* 2020;392:112356.
40. Pejović A, Minić A, Bugarinović J, Pešić M, Damjanović I, Stevanović D, et al. Synthesis, characterization and antimicrobial activity of novel 3-ferrocenyl-2-pyrazolyl-1, 3-thiazolidin-4-ones. *Polyhedron* 2018;155:382-9.
41. Ismail AM, El Maaty WM, Jean-Claude BJ, Mostafa SI. Synthesis, characterization and anticancer activity of new Zn (II) and MoO<sub>2</sub>+ complexes of 2-amino-4, 6-mercaptotriazine. *Inorg Chem Commun* 2019;106:217-23.
42. Moteki H, Hibasami H, Yamada Y, Katsuzaki H, Imai K, Komiya TA. Specific induction of apoptosis by 1, 8-cineole in two human leukemia cell lines, but not in human stomach cancer cell line. *Oncol Rep* 2002;9:757-60.
43. Cowley AR, Davis J, Dilworth JR, Donnelly PS, Dobson R, Nightingale A, et al. Fluorescence studies of the intra-cellular distribution of zinc bis(thiosemicarbazone) complexes in human cancer cells. *Chem Commun* 2005;7:845-7.